

IMAGEM EM NEUROLOGIA/IMAGE IN NEUROLOGY

Let the Fibers Guide You: DTI in ARSACS

Deixa as Fibras Guiarem-te: DTI na ARSACS

 José Sá Silva ^{1,*},  Miguel Mendonça Pinto ²,  Catarina Mendes Pinto ¹

1-Serviço de Neurorradiologia, Centro Hospitalar Universitário do Porto, Porto, Portugal

2-Unidade de Neuropatologia, Centro Hospitalar Universitário do Porto, Porto, Portugal

DOI: <https://doi.org/10.46531/sinapse/IN/210083/2022>

A 35-year-old Portuguese woman with a history of parental consanguinity, but no family history of neuropathy, was followed from the age of 9 years for a demyelinating peripheral neuropathy. Initial nerve conduction studies and electromyography revealed slowed conduction velocity and low amplitude motor and sensory responses, especially in lower extremities. With disease progression, Charcot-Marie-Tooth (CMT) disease was suspected and a next generation sequencing (NGS) based gene panel test for CMT type 4 was used (*EGR2*, *FGD4*, *FIG4*, *GDAPI*, *HKI*, *MTMR2*, *NDRG1*, *PRX*, *SBF1*, *SBF2* and *SH3TC2*) – although no abnormal variant was identified, CMT remained as the suspected diagnosis.

The clinical picture was initially dominated by distal motor weakness, predominantly of the lower limbs. Over the course of several years it progressed to include cerebellar upper-limb ataxia with oculomotor disturbances and dysarthria, lower-limb spasticity, distal muscle atrophy and weakness, absent reflexes, and impaired proprioception – these were all findings characteristic of autosomal recessive spastic ataxia of Charlevoix-Saguenay (ARSACS).¹ There was no cognitive impairment. It was the onset of cerebellar findings that raised suspicion of a missed diagnosis and prompted a NGS gene panel test for hereditary neuropathies, which uncovered a homozygous pathogenic variant of SACS, a gene located on chromosome 13q which encodes the sascin protein.² Further investigation with magnetic resonance imaging

(MRI) revealed typical imaging findings – atrophy of superior vermis, a bulky pons with bilateral, parallel, paramedian, linear hypointensities in T2 weighted-images and linear T2-hyperintensities surrounding the lateral thalami (Fig. 1). DTI further confirmed the diagnosis, revealing bulky transverse pontine fibers (red in Fig. 1) and a thin cortico-spinal tract (blue in Fig. 1).




Figure 1. MRI revealed atrophy of the superior cerebellar vermis (A, long-arrow), a bulky pons (A, short-arrow) with paramedian T2-hypointensity (B), thinning of corpus callosum posterior midbody (A, arrowhead) and linear T2-hyperintensity around the thalami (C). DTI showed over-represented transverse pontine fibers (D, arrow, in red) displacing the abnormally small cortico-spinal tract (D, arrowhead, in blue).

Although the classical findings in ARSACS are well described on conventional MRI,³ findings in advanced MRI techniques, specifically in diffusion tensor imaging (DTI), are not so well established. Nevertheless, DTI has been increasingly pointed as a very specific technique to diagnose ARSACS patients, with very distinct

Informações/Informations:

Imagem em Neurologia, publicado em Sinapse, Volume 22, Número 2, abril-junho 2022. Versão eletrónica em www.sinapse.pt; Image in Neurology, published in Sinapse, Volume 22, Number 2, April-June 2022. Electronic version in www.sinapse.pt
 © Autor (es) (ou seu (s) empregador (es)) e Sinapse 2022. Reutilização permitida de acordo com CC BY-NC. Nenhuma reutilização comercial. © Author(s) (or their employer(s)) and Sinapse 2022. Re-use permitted under CC BY-NC. No commercial re-use.

Keywords:

Muscle Spasticity; Spinocerebellar Ataxias.

Palavras-chave:

Ataxias Espinocerebelares; Espasticidade Muscular.

***Autor Correspondente / Corresponding Author:**

José Sá Silva
 Serviço de Neurorradiologia, Centro Hospitalar Universitário do Porto
 Largo do Prof. Abel Salazar, 4099-001 Porto, Portugal
jose.msas.silva@gmail.com

Recebido / Received: 2021-12-20

Aceite / Accepted: 2022-05-07

Publicado / Published: 2022-06-30

tive features.⁴ These features include a relative excess of transverse pontine fibers (which appear thickened) and small cortico-spinal tracts.⁴ The combination of classical findings in conventional MRI might be relatively specific of ARSACS, but those findings can be variably present and more or less evident,³ making DTI a useful confirmatory tool when conventional MRI is uncertain.

Clinically, ARSACS can have different phenotypes and the clinical presentation can take years to be fully expressed, which leads to underdiagnosis of ARSACS. This case highlights the possible misleading presenting phenotype of pure neuropathy, in particular resembling Charcot-Marie-Tooth disease,¹ emphasizing the need for sensitive diagnostic tools. Although not widely used, DTI can be a very good, non-invasive, and highly specific tool to diagnose this disease. ■

Contributorship Statement / Declaração de Contribuição

JSS: Conception and writing of the manuscript and preparation of the images.

MMP: Data interpretation and critical review.

CMP: Writing of the manuscript and critical review.

Responsabilidades Éticas

Conflitos de Interesse: Os autores declaram a inexistência de conflitos de interesse na realização do presente trabalho.

Fontes de Financiamento: Não existiram fontes externas de financiamento para a realização deste artigo.

Confidencialidade dos Dados: Os autores declaram ter seguido os protocolos da sua instituição acerca da publicação dos dados de doentes.

Consentimento: Consentimento do doente para publicação obtido.

Proveniência e Revisão por Pares: Não comissionado; revisão externa por pares.

Ethical Disclosures

Conflicts of Interest: The authors have no conflicts of interest to declare.

Financing Support: This work has not received any contribution, grant or scholarship.

Confidentiality of Data: The authors declare that they have followed the protocols of their work center on the publication of data from patients.

Patient Consent: Consent for publication was obtained.

Provenance and Peer Review: Not commissioned; externally peer reviewed.

References / Referências

1. Synofzik M, Soehn AS, Gburek-Augustat J, Schicks J, Karle KN, Schüle R, et al. Autosomal recessive spastic ataxia of Charlevoix Saguenay (ARSACS): expanding the genetic, clinical and imaging spectrum. *Orphanet J Rare Dis*. 2013;8:41. doi: 10.1186/1750-1172-8-41.
2. Vermeer S, van de Warrenburg BP, Kamsteeg EJ, Brais B, Synofzik M. ARSACS. 2003 Dec 9 [Updated 2020 Jan 2]. In: Adam MP, Ardinger HH, Pagon RA, editors. *GeneReviews*® [Internet]. Seattle: University of Washington, Seattle; 1993-2022. [accessed Oct 2021] Available from: <https://www.ncbi.nlm.nih.gov/books/NBK1255/>
3. Prodi E, Grisoli M, Panzeri M, Minati L, Fattori F, Erbetta A, et al. Supratentorial and pontine MRI abnormalities characterize recessive spastic ataxia of Charlevoix-Saguenay. A comprehensive study of an Italian series. *Eur J Neurol*. 2013;20:138-46. doi: 10.1111/j.1468-1331.2012.03815.x.
4. Oguz KK, Haliloglu G, Temucin C, Gocmen R, Has AC, Dörschner K, et al. Assessment of whole-brain white matter by DTI in autosomal recessive spastic ataxia of Charlevoix-Saguenay. *AJNR Am J Neuroradiol*. 2013;34:1952-7. doi: 10.3174/ajnr.A3488.